



LARGE ROTATOR CUFF REPAIR TEAR >3 cm

ROM GOALS

	PFE	PER@20	PER@90	AFE
POW 2	60°-90°	0°-20°	NA	NA
POW 6	90°-120°	20°-30°	NA	To 90°
POW 9	130°-155°	30°-45°	45°-60°	90°-120°
POW 12	140°-WNL	30°-WNL	75°-WNL	120°-WNL

Phase I: (Protective Phase) – NO PROM UNTIL POW 2

Goals: Minimize pain and inflammatory response
Achieve ROM goals
Establish stable scapula

Weeks 0 to 6

1. Elbow, wrist and hand AROM (EWH)
2. Supine passive external rotation (PER)
 - a. T-stick in 0-20 deg flexion and 20 deg abduction
 - b. 10 reps, 2 x day
3. Supine passive forward elevation in plane of scapula
 - a. 90 after the first 2 weeks
 - b. 10-20 reps, 2 x day
 - c. **Do not begin passive forward elevation until 2 weeks post op**
4. C-spine AROM
5. Ice
6. Positioning full time in sling with abduction pillow
7. Shoulder shrugs and retractions (no weight)
8. ***Pain control modalities PRN
9. Complications/Cautions:
 - a. If pain level is not dissipating, decrease intensity and volume of exercises.

- b. Assure normal neurovascular status
- c. No AAROM or AROM until 6 weeks**
- d. No pulley until 6 weeks**

Weeks 6 to 12

1. Heat/ice PRN to help obtain motion
2. D/C sling as comfortable
3. Progress PROM goals
4. Achieve PROM goals in ER at 20 degrees and 90 degrees abduction
- 5. Start AAFE and progress to AFE**
6. Initiate posterior capsule stretching
7. Isometrics, keeping elbow flexed to 90 degrees (Sub maximal, pain free)
8. Theraband scapula retractions
9. ***Aquatics
- 10.***Mobilizations PRN
- 11.***Trunk stabilization/strengthening
- 12.Start periscapular strengthening with very low weight and high repetitions
- 13.Cautions:
 - a. Do not initiate rotator cuff strengthening until 12 weeks and until night pain has subsided and overall pain level is low**

Phase II: (Progressive Strengthening)

Goals: Achieve staged ROM goals
Eliminate shoulder pain
Improve strength, endurance and power
Increase functional activities

Months 3 to 4

1. Continue as above
2. ROM should be full in all planes
3. Progress isometrics
4. Advance scapula strengthening
5. ***Mobilizations PRN
6. ***Aquatics for strengthening
7. ***CKC activities for dynamic stability of scapula deltoid and cuff
8. ***Trunk stabilization/strengthening
9. ***Light PNF D1, D2 and manual resistance for cuff/deltoid/scapula (rhythmic stabilization or slow reversal hold)

10. Initiate theraband ER and IR strengthening
11. Progressive serratus anterior strengthening
(isolated pain free, elbow by side)
12. Progress to isotonic dumbbell exercises for deltoid, supraspinatus, up to 3 lbs max

13. Cautions:

- a. Do not initiate AAFE or theraband rotator cuff strengthening until overall pain level is low
- b. Assure normal scapulohumeral rhythm with AAFE and AFE
- c. Strengthening program should progress only without signs of increasing inflammation
- d. Strengthening program should emphasize high repetitions, low weight and should be performed a maximum of 2 x day

Phase III: (Return to activity/advanced conditioning)

Goals: Normalize strength, endurance and power
Return to full ADL's and recreational activities

Month 4 to 6

1. Stretching PRN
2. Continue deltoid/cuff/scapula strengthening as above (5lbs max for isotonic strengthening) with the following progressions:
 - a. Prone isotonic strengthening PRN
 - b. Decreasing amounts of external stabilization provided to shoulder girdle
 - c. Integrate functional patterns
 - d. Increase speed of movements
 - e. Integrate kinesthetic awareness drills into strengthening activities
 - f. Decrease in rest time to improve endurance
3. May begin tennis ground stroke/batting/return to golf after completing strengthening progression
4. ***Progressive CKC dynamic stability activities
5. ***Impulse
6. ***Initiate isokinetic strengthening
7. ***Mobilizations PRN
8. ***Trunk stabilization/strengthening

Month 6 to 8

1. Stretching PRN
2. Continue deltoid/cuff/scapula strengthening program
3. Initiate plyometric program (if needed)
 - a. Do not begin until 5/5 MMT for rotator cuff and scapula
 - b. QD at most
 - c. Begin with beach ball/tennis ball progressing to weighted balls
 - d. 2-handed tosses: waist level
 - i. Overhead
 - ii. Diagonal
 - e. 1-handed stability drills
 - f. 1-handed tosses (vary amount of abduction, UE support, amount of protected ER)
4. May begin Interval Throwing Program after 3-6 weeks of plyometrics
5. Initiate progressive replication of demanding ADL/work activities

Discharge/Return to sport criteria

1. PROM WNL for ADL's/work/sports
2. MMT 5/5 shoulder girdle and/or satisfactory isokinetic test
3. Complete plyometric program, if applicable
4. Complete interval return to sport program, if applicable