

**OPEN SHOULDER RECONSTRUCTION OR LATERJET PROCEDURE
 REHABILITATION PROTOCOL**

	RANGE OF MOTION*	IMMOBILIZER	THERAPEUTIC EXERCISE
PHASE I 3 - 6 weeks	Active/Active-Assistive: stretch to 20° of external rotation, and 90° of forward flexion-internal rotation to stomach Abduction to 45 degrees No cross-body adduction Protect the subscap!!!	Worn at all times except for hygiene and therapeutic exercise	Elbow/wrist/hand ROM, grip strengthening, isometric in sling, external/internal rotation exercises with elbow at side
PHASE II 6 - 12 weeks	Increase forward flexion to 140/40° ER at side/ 60° abduction/ IR behind back to waist	None	Advance isometrics in phase I to use of a theraband, continue with elbow/wrist/hand ROM and grip strengthening, begin prone extensions, and scapular stabilizing exercises, gentle joint mobs Modalities as needed
PHASE III 12 - 16 weeks	Progress to full active motion without discomfort Gentle passive stretching at end ranges	None	Advance theraband exercises to use of weights, continue with and progress exercises in phase II Include RTC, deltoid, and scapular stabilizers
PHASE IV	Full without discomfort	None	Advance exercises in phase III, begin functional progression to work/sport, return to previous activity level* begin upper body ergometer, proprioception plyometrics

*Patient may return to the weight room at 3 months, if appropriate

**Patient may return to competitive sports, including contact sports, by 6 months, if approved