

# CALF STRAIN

## What is a calf strain?

A strain is an injury in which muscle fibers or tendons are stretched or torn. People commonly call such an injury a “pulled” muscle. A calf strain is an injury to the muscles and tendons in the back of your leg below your knee.

## How does it occur?

A strain of your calf muscles can occur during a physical activity where you push off forcefully from your toes. It may occur in running, jumping, or lunging.

## What are the symptoms?

A calf muscle strain may cause immediate pain in the back of your lower leg. You may hear or feel a pop or a snap.

You may get the feeling that someone has hit you in the back of the leg. It is hard to rise up on your toes. Your calf may be swollen and bruised.

## How is it diagnosed?

Your healthcare provider will examine your lower leg. Your calf muscles will be tender.

## How is it treated?

Treatment may include:

- applying ice packs to your calf for 20 to 30 minutes every 3 to 4 hours for 2 or 3 days or until the pain goes away
- elevating your leg on a pillow while you are lying down
- wrapping an elastic bandage around your calf to keep the swelling from getting worse
- using crutches, if it is too painful to walk.
- taking anti-inflammatory medicine (adults aged 65 years and older should not take non-steroidal anti-inflammatory medicine for more than 7 days without their healthcare provider’s approval)
- getting physical therapy, which may include treatment of the muscle tissue by a therapist using ultrasound or muscle stimulation.
- having your healthcare provider or therapist tape the injured muscles while they are healing to help you to return to athletic activities
- doing rehabilitation exercises

While you are recovering from your injury, you will need to change your sport or activity to one that does not make your condition worse. For example, you may need to swim instead of run.


## When can I return to my sport or activity?

The goal of rehabilitation is to return you to your sport or activity as soon as is safely possible. If you return too soon you may worsen your injury, which could lead to permanent damage. Everyone recovers from injury at a different rate. Return to your activity will be determined by how soon your calf recovers, not by how many days or weeks it has been since your injury occurred. In general, the longer you have symptoms before you start treatment, the longer it will take to get better.

You may safely return to your sport or activity when, starting from the top of the list and progressing to the end, each of the following is true:

- You have full range of motion in the injured leg compared to the uninjured leg.
- You have full strength of the injured leg compared to the uninjured leg.
- You can jog straight ahead without pain or limping.

## CALF STRAIN



- You can sprint straight ahead without pain or limping.
- You can do 45-degree cuts, first at half-speed, then at full-speed.
- You can do 20-yard figures-of-eight, first at half-speed, then at full-speed.
- You can do 90-degree cuts, first at half-speed, then at full-speed.
- You can do 10-yard figures-of-eight, first at half-speed, then at full-speed.

- You can jump on both legs without pain and you can jump on the injured leg without pain.


### How can calf strains be prevented?

Calf strains are best prevented by warming up properly and doing calf-stretching exercises before your activity. This is especially important if you are doing jumping or sprinting sports.

## CALF STRAIN REHABILITATION EXERCISES


You can begin gently stretching your calf muscle using the towel stretch right away. Make sure you only get a gentle pull and not a sharp pain while you are doing this stretch.

**1. TOWEL STRETCH:** Sit on a hard surface with one leg stretched out in front of you. Loop a towel around the ball of your foot and pull the towel toward your body keeping your knee straight. Hold this position for 15 to 30 seconds then relax. Repeat 3 times.



TOWEL STRETCH


**3. RESISTED ANKLE PLANTAR FLEXION:** Sit with your leg outstretched and loop the middle section of the tubing around the ball of your foot. Hold the ends of the tubing in both hands. Gently press the ball of your foot down and point your toes, stretching the tubing. Return to the starting position. Do 3 sets of 10.



RESISTED ANKLE PLANTAR FLEXION

After you can do the towel stretch easily, you can start the standing calf stretch.

**2. STANDING CALF STRETCH:** Facing a wall, put your hands against the wall at about eye level. Keep one leg back with the heel on the floor, and the other leg forward. Turn your back foot slightly inward (as if you were pigeon-toed) as you slowly lean into the wall until you feel a stretch in the back of your calf. Hold for 15 to 30 seconds. Repeat 3 times. Do this exercise several times each day.



STANDING CALF STRETCH

After a couple days of stretching, you can begin strengthening your calf and lower leg muscles using elastic tubing as described in the next exercise.

You may do the last 4 exercises when you can stand on your toes without pain.

**4. HEEL RAISE:** Balance yourself while standing behind a chair or counter. Raise your body up onto your toes and hold for 5 seconds. Then slowly lower yourself down. Hold onto the chair or counter if you need to. When this exercise becomes less painful, try lowering on one leg only. Repeat 10 times. Do 3 sets of 10.



HEEL RAISE

You can challenge yourself by standing only on your injured leg and lifting your heel off the ground.



**5. SINGLE LEG BALANCE:** Stand without any support and attempt to balance on one leg. Begin with your eyes open and then try to perform the exercise with your eyes closed. Hold the single-leg position for 30 seconds. Repeat 3 times. When you have mastered this, try doing this exercise standing on a pillow.

SINGLE LEG BALANCE


**6. NOSE TOUCH:** Stand on one leg facing a wall. Stand 4 inches from the wall. Keep your body and leg straight. Slowly lean forward, trying to touch your nose to the wall. Make sure you do not bend forward at your waist. Do 3 sets or 10.



NOSE TOUCH

**7. WALL JUMP:** Face a wall and place a piece of masking tape about 2 feet above your head. Jump up with your arms above your head and try to touch the piece of tape. Make sure you do a “spring” type of motion and do not land hard onto your feet. Progress to taking off and landing on one foot. Do 3 sets of 10.

Another good exercise is hopping. You can start at one end of the room and try to hop as high as you can across the room on one foot. Jumping rope is also a good exercise.



WALL JUMP