NECK STRAIN

What is a neck strain?

A strain is a tear of a muscle or tendon. Your neck is surrounded by small muscles, that run close to the vertebrae, and larger muscles, that make up the visible muscles of the neck.

How does it occur?

Neck strains most often occur, when the head and neck are forcibly moved, such as in a whiplash injury or from contact in sports. Sometimes strains can occur from an awkward position during sleep or poor posture while working at a computer.

What are the symptoms?

You have pain in your neck. When the neck muscles go into spasm you feel hard, tight muscles in your neck that are very tender to the touch. You have pain when you move your head to either side or when you try to move your head up or down. The spasming muscles can cause headaches.

The pain may start right after an injury or may take a few hours or days to develop. Other symptoms may include neck stiffness, dizziness, or unusual sensations, such as burning or a pins-and-needles feeling.

How is it diagnosed?

Your healthcare provider will examine your neck. Your neck muscles will be tender and tight. You may have pain over the bones in your neck. Your healthcare provider may order X-rays to make sure the vertebrae are not injured.

How is it treated?

Right after the injury you should place an ice pack on your neck for 20 to 30 minutes every 3 or 4 hours for 2 to 3 days or until the pain goes away.

Your healthcare provider may prescribe an anti-inflammatory medication and a neck collar to support your neck and prevent further injury (adults aged 65 years and older should not take non-steroidal anti-inflammatory medicine for more than 7 days without their healthcare provider's approval).

If you still have neck pain several days after the injury and after using ice, your healthcare provider may recommend using moist heat on your neck. You can buy a moist-heat pad or make your own by soaking towels in hot water. Put moist heat on your neck for 20 to 30 minutes every 3 or 4 hours

until the pain goes away. You may find that it helps to alternate putting heat and ice on your neck.

How long will the effects last?

The length of recovery depends on many factors such as your age, health, and if you have had a previous neck injury. Recovery time also depends on the severity of the injury. A mild injury may recover within a few weeks, whereas a severe injury may take 6 weeks or longer to recover. Ask your healthcare provider when you can return to your normal activities.

When can I return to my sport or activity?

The goal of rehabilitation is to return you to your sport or activity as soon as is safely possible. If you return too soon you may worsen your injury, which could lead to permanent damage. Everyone recovers from injury at a different rate. Return to your sport or activity will be determined by how soon your neck recovers, not by how many days or weeks it has been since your injury occurred. In general, the longer you have symptoms before you start treatment, the longer it will take to get better.

If you participate in contact sports, it is important to rehabilitate your neck and shoulders before going back to competition. You must have full range of motion of your neck. This means you must be able to:

NECK STRAIN



PAGE 1 OF 3 PAGES

8. ISOMETRIC NECK EXTENSION: Sit tall, eyes straight ahead, and chin level. Clasp your hands together and place them behind your head. Press the back of your head into your palms. Hold 5 seconds and release.

Do 3 sets of 5. ISOMETRIC NECK EXTENSION




ISOMETRIC NECK SIDE BEND

9. ISOMETRIC NECK SIDE BEND: Sit tall, eyes straight ahead, and chin level. Place the palm of your hand at the side of your temple and press your temple into the palm of your hand. Hold 5 seconds and release. Do 3 sets of 5 on each side.

10. SCAPULAR SQUEEZE: While sitting or standing with your arms by your sides, squeeze your shoulder blades together and hold for 5 seconds. Do 3 sets of 10.

SCAPULAR SQUEEZE



12. HEAD LIFT: Neck curl: Lie on your back with your knees bent and your feet flat on the floor. Tuck your chin and lift your head toward your chest, keeping your shoulders on the

5 seconds. Repeat 10 times.

floor. Hold for

HEAD LIFT: NECK CURL

13. HEAD LIFT: NECK SIDE BEND: Lie on your right side with your right arm laying straight out. Rest your head on your arm, then lift your head slowly toward your left shoulder. Hold for 5 seconds. Repeat 10 times. Switch to your left side and repeat the exercise lifting your head toward your right shoulder.

HEAD LIFT: NECK SIDE BEND

 $Copyright @ 2010 \ McKesson \ Corporation \ and /or \ its \ affiliates. \ All \ rights \ reserved. \ Permission \ to \ copy \ for \ patient \ education. \ www.sportsmedpress.com \ and \ for \ reserved.$